

On the divergences of inflationary superhorizon perturbations

K. Enqvist^{1,2}, S. Nurmi^{1,2}, D. Podolsky², and G. I. Rigopoulos²
¹*Physics, Department, University of Helsinki; ²Helsinki Institute of Physics,*
P.O.Box 64, FIN-00014, University of Helsinki, Finland

We discuss the infrared divergences that appear to plague cosmological perturbation theory. We show that within the stochastic framework they are regulated by eternal inflation so that the theory predicts finite fluctuations. Using the ΔN formalism to one loop, we demonstrate that the infrared modes can be absorbed into additive constants and the coefficients of the diagrammatic expansion for the connected parts of two and three-point functions of the curvature perturbation. As a result, the use of any infrared cutoff below the scale of eternal inflation is permitted, provided that the background fields are appropriately redefined. The natural choice for the infrared cutoff would of course be the present horizon; other choices manifest themselves in the running of the correlators. We also demonstrate that it is possible to define observables that are renormalization group invariant. As an example, we derive a non-perturbative, infrared finite and renormalization point independent relation between the two-point correlators of the curvature perturbation for the case of the free single field.

I. INTRODUCTION

Primordial perturbations generated during inflation are conveniently characterized by the gauge-invariant curvature perturbation which can be directly related to the observed CMB anisotropies [1]. The study of the properties of the curvature perturbation arising in various inflationary models has led to a remarkable confrontation of early universe theory with cosmological observations. The increasing observational precision has placed increasing demands on the theory and over the past few years results beyond linear theory have been sought, mainly focusing, but not limited to, non-gaussianity. It is well known however that the n -point correlators of the curvature perturbation contain IR divergences as a consequence of the approximative scale invariance and gaussianity of inflationary perturbations¹. Although correlators extending to superhorizon scales as such are not observable, the infrared behavior can manifest itself non-trivially both in classical and quantum field theory when computing higher order perturbative corrections. Moreover, in principle we only have to wait long enough, and the infrared modes will become observable. Therefore, a satisfactory understanding of the theory requires that these IR divergences can be controlled.

To demonstrate the problems related to the infrared divergences, it is enough to consider two point correlators and single field inflation. Using the ΔN formalism [2], the curvature perturbation $\zeta(\mathbf{x})$ can be expressed in a simple form

$$\zeta(\mathbf{x}) = N'\phi(\mathbf{x}) + \frac{1}{2}N''(\phi^2(\mathbf{x}) - \langle\phi^2\rangle) + \dots, \quad (1)$$

where N is the number of e-folds and ϕ denotes perturbations of the inflaton field on spatially flat slices. The first order contribution to the two point correlator of the curvature perturbation behaves as

$$\langle\zeta(\mathbf{x}_1)\zeta(\mathbf{x}_2)\rangle_{(1)} = (N')^2 G(|\mathbf{x}_1 - \mathbf{x}_2|) \sim -(N')^2 \mathcal{P}_\phi \ln\left(\frac{|\mathbf{x}_1 - \mathbf{x}_2|}{L}\right), \quad (2)$$

where $G(|\mathbf{x}_1 - \mathbf{x}_2|) = \langle\phi(\mathbf{x}_1)\phi(\mathbf{x}_2)\rangle$ is the correlation function of the scalar field, $\mathcal{P}$ gives the amplitude of the inflaton perturbations ϕ , and we have considered the case of scale invariant fluctuations. The scale L is an a priori arbitrary cutoff needed to regulate the infrared divergences of the two point correlator. In reality, the theory may contain a physical scale beyond which scale invariance is broken and L may be related to that scale. For instance, one could argue that in our local patch inflation has lasted only a finite time so that beyond some infrared scale, the assumption of scale invariance necessarily breaks down. In any case, in the actual experiments only differences of correlators are measured and they remain finite even in the limit $L \rightarrow \infty$,

$$\langle\zeta(\mathbf{x}_1)\zeta(\mathbf{x}_2)\rangle_{(1)} - \langle\zeta(\mathbf{x}_1)\zeta(\mathbf{x}_3)\rangle_{(1)} \sim -(N')^2 \mathcal{P}_\phi \ln\left|\frac{\mathbf{x}_1 - \mathbf{x}_2}{\mathbf{x}_1 - \mathbf{x}_3}\right|. \quad (3)$$

The value of the IR cutoff L is therefore irrelevant when considering observable quantities calculated using first order perturbation theory.

¹ More accurately, the expressions depend on an IR cutoff L .

The situation becomes more complicated beyond the linear order since the divergent part of the correlators will no longer be simply an additive constant. For example, the second order contribution to the two point correlator behaves as

$$\langle \zeta(\mathbf{x}_1)\zeta(\mathbf{x}_2) \rangle_{(2)} \sim -N'N''' \langle \phi^2 \rangle \mathcal{P}_\phi \ln\left(\frac{|\mathbf{x}_1 - \mathbf{x}_2|}{L}\right) + \frac{1}{2}(N'')^2 \mathcal{P}^2 \ln^2\left(\frac{|\mathbf{x}_1 - \mathbf{x}_2|}{L}\right), \quad (4)$$

which in the limit $L \rightarrow \infty$ contains divergences depending also on the separation of the two points $\mathbf{x}_1$ and $\mathbf{x}_2$.

There are two distinct issues associated with the appearance of these IR divergences. The first is whether $G(|\mathbf{x}_1 - \mathbf{x}_2|)$ is actually IR divergent. The divergence is the outcome of the assumption that the spectrum is scale invariant, an assumption that will probably break down after some scale L . One might expect that the complete theory of inflation predicts a well defined value for $G(|\mathbf{x}_1 - \mathbf{x}_2|)$, maybe large but free of divergences. The other issue is the relation of this (large) value of $G(|\mathbf{x}_1 - \mathbf{x}_2|)$ to the measurements of an observer accessing only a small patch of the whole universe. In other words: how do the predictions of the theory defined on the largest possible scales translate into predictions for measurements on much smaller scales? These problems have been discussed in the literature [3, 4, 5, 6, 7], but there seems to be no agreement on the correct approach.

In this paper we address both issues. First, we show that the variance of the field $\langle \phi^2 \rangle$ is finite and well defined for generic slow-roll inflation. We prove this using the framework of stochastic inflation which accurately describes the fluctuations of quantum fields on superhorizon scales. Furthermore, we note that since IR fluctuations are only observationally relevant in regions which have exited the eternally inflating regime and have eventually thermalized, the scale of eternal inflation $V(\phi_{\text{EI}}) \lesssim \epsilon(\phi_{\text{EI}})M_P^4$ provides an ultimate upper limit for observable fluctuations. Only below the length scale L_{EI} , corresponding to ϕ_{EI} , does the notion of a well defined evolving background field apply and the spacetime can be approximated by a global FRW metric. We then proceed to consider how predictions at this scale can be related to predictions for observations restricted in patches of size $M < L < L_{\text{EI}}$ by applying a systematic renormalization prescription. In particular, we focus on the renormalization of the classical one loop expressions for the 2-point and 3-point functions of the curvature perturbation. We find that the renormalization amounts to shifting the background fields in the connected parts of the correlators. Furthermore, we note that apart from some contrived models, the exact position of the patch of size M in the larger universe is irrelevant in single field inflation, to the extent that the background evolution corresponding to the observable patch M has been defined. Therefore, averaging over all possible embeddings of M in the whole universe has no meaning in this context.

The paper is organized as follows: In sections II and III we use the formalism of stochastic inflation to demonstrate that for potentials supporting slow-roll inflation, the one point functions of the scalar field and the curvature perturbation are finite as long as the energies are sub-planckian. Therefore there are no real IR divergences associated with inflationary fluctuations. Then, in section IV we relate predictions at scales L to scales $M < L$ using the ΔN formalism, and argue that a renormalization group equation can be used to relate predictions obtained using two different cutoffs/renormalization scales. In section V we discuss a concrete example, the free field, and demonstrate that it is possible to construct an observable that is both infrared finite and renormalization point independent. In section VI we close with a discussion and a comparison of our findings to recent work on the subject.

II. ETERNAL INFLATION AND ONE-POINT CORRELATION FUNCTIONS OF THE INFLATON

In this Section, we will show how effects of eternal inflation lead to the regularization of the infrared divergences present in the correlation functions of the inflaton. Although we will focus on the chaotic inflation scenario [8, 9], our analysis can be easily extended to the case of new inflation [8, 10], inflation driven by multiple fields [11] or to the case of the curvaton scenario [12].

Let us suppose that primordial inflation is driven by a single scalar field ϕ with the potential $V(\phi)$ satisfying the slow roll initial conditions $\epsilon = \frac{1}{2}M_P^2 (V_{,\phi}/V)^2 \ll 1$, $\eta = M_P^2 V_{,\phi\phi}/V \ll 1$. During one Hubble time $\Delta t \sim H^{-1}$ the value of the inflaton field changes by

$$\Delta\phi \sim \dot{\phi}\Delta t \sim \frac{V_{,\phi}}{3H^2} \sim \frac{M_P^2}{8\pi} \frac{V_{,\phi}}{V}. \quad (5)$$

On the other hand, at the same time scale fluctuations of the inflaton $\delta\phi$ are generated with a characteristic wavelength $l \sim k^{-1} \sim H^{-1}$. These fluctuations have a randomly distributed amplitude among different causally disconnected regions (Hubble patches), and the width of this distribution is given by

$$|\delta\phi| \sim \frac{H}{2\pi} \sim \sqrt{\frac{2V}{3\pi M_P^2}}. \quad (6)$$

As a result, observers living in different Hubble patches see different expectation values of the inflaton, because for them long wavelength (superhorizon) fluctuations $\delta\phi$ are physically indistinguishable from the inflaton zero mode.

In some Hubble patches the sign of the infrared fluctuation $\delta\phi$ is positive, in others it is negative and therefore may overcome the effect of the classical force (5) acting on the inflaton. When $\Delta\phi \lesssim |\delta\phi|$, stochastic fluctuation $\delta\phi$ may lead to the effective growth of the expectation value of the inflaton ϕ in a given Hubble volume. This regime is denoted as *eternal inflation* because there will be always Hubble patches where the expectation value of the inflaton continues to grow [13].

One can see that the stochastic force becomes more important than the classical one when $|(M_P^2/V)V_{,\phi}| \lesssim \sqrt{V/M_P^2}$ or

$$V(\phi_{\text{EI}}) \gtrsim \epsilon(\phi_{\text{EI}})M_P^4, \quad (7)$$

where ϵ is the slow roll parameter. For example, in the $\lambda\phi^4$ chaotic inflation model the condition (7) is satisfied when $\phi > \phi_{\text{EI}} = \lambda^{-1/6}M_P$. Since the upper bound on possible values of field corresponds to the Planckian energy density, we conclude that the regime of eternal inflation is realized at $\lambda^{-1/6}M_P < \phi < \lambda^{-1/4}M_P$, a very wide interval of possible values of the inflaton, since $\lambda \sim 10^{-13}$.

Larger values of field correspond to larger scales of perturbations in the chaotic inflationary scenario, and one may conclude that the regime of eternal inflation can influence the infrared behavior of the correlation functions of both ϕ and the curvature perturbation ζ . In fact, as Woodard has shown [14], leading infrared logarithmic divergences in the correlation functions of ϕ and ζ can be reproduced in the formalism of stochastic inflation [8] which is thus well suited for treating the regime of eternal inflation.

In this formalism one decomposes the Heisenberg operator for the field ϕ into the infrared and ultraviolet parts according to the prescription

$$\hat{\phi}(N, \mathbf{x}) = \Phi(N, \mathbf{x}) + \frac{1}{(2\pi)^{3/2}} \int d^3k \theta(k - \delta aH) \left(\hat{a}_k \phi_k(N) e^{-i\mathbf{k}\mathbf{x}} + \hat{a}_k^\dagger \phi_k^*(N) e^{i\mathbf{k}\mathbf{x}} \right) + \delta\phi, \quad (8)$$

where $N = \log a$ is the number of inflationary e-folds, $\delta \ll 1$ is a small parameter², $\theta(x)$ is the Heaviside step function and $\delta\phi$ is a small contribution suppressed by additional powers of the slow roll parameters. We are especially interested in the dynamics of the infrared part Φ of the inflaton field, which is responsible for the infrared divergences.

Substituting the decomposition (8) into the equation of motion for the Heisenberg operator $\hat{\phi}$, one can show that the infrared part Φ satisfies the Langevin equation

$$\frac{\partial \Phi}{\partial N} = -\frac{1}{3H^2} V_{,\Phi} + \frac{f}{H}, \quad (9)$$

where f is the composite operator containing contributions of the ultraviolet modes and having the correlation property

$$\langle f(N) f(N') \rangle = \frac{H^4}{4\pi^2} \delta(N - N'), \quad (10)$$

as can be checked by a direct calculation of this correlator in the Bunch-Davies vacuum state. The remarkable property of the equation (9) is that all its terms commute with each other and therefore can be considered as classical stochastic quantities. The Langevin equation (9) describes the process of the inflaton random walk due to the effect of generated superhorizon fluctuations.

At the next step, by the Stratonovich prescription one proceeds from the Langevin equation (9) to the Fokker-Planck equation

$$\frac{\partial P}{\partial N} = -\frac{1}{3\pi M_P^2} \frac{\partial^2}{\partial \phi^2} (VP) + \frac{M_P^2}{8\pi} \frac{\partial}{\partial \phi} \left(V^{-1} \frac{\partial V}{\partial \phi} P \right), \quad (11)$$

which governs the evolution of the probability $P(\phi, N)$ to measure a given expectation value of the inflation in a given Hubble patch. Its solution can be expressed in the form

$$P(\phi, N) = v(\phi) e^{-v(\phi)} \sum_{n=0}^{+\infty} c_n \psi_n(\phi) e^{-E_n N}, \quad (12)$$

² This parameter is intrinsically related to the decoherence scale l_d , where perturbations which are leaving the horizon finally become classical (i.e., to the scale, where decaying mode can be neglected as compared to the growing one). The scale decoherence l_d corresponds to few e-folds after perturbations leave the horizon. Note that δ cannot be made arbitrarily small as it is related to the slow roll parameters [8].

where

$$v(\phi) = \frac{3M_P^4}{16V(\phi)}, \quad (13)$$

while ψ_n and E_n are the eigenfunctions and eigenvalues of the Schrödinger equation given by

$$-\frac{1}{2}\psi_n'' + \frac{1}{2}(-v'' + (v')^2)\psi_n = E_n\psi_n \frac{8\pi v}{M_P^2}. \quad (14)$$

Due to the supersymmetric form of the potential in the Schrödinger equation (14) all eigenvalues satisfy the condition $E_n \geq 0$, and it is trivial to see that the ground state corresponds to a zero eigenvalue. To be precise, the ground state corresponds to $E_0 = 0$ if and only if the corresponding eigenmode is normalizable. However, this is always the case for chaotic inflationary models with $V(\phi)$ bounded from below and growing as $\phi \rightarrow 0$.

The Fokker-Planck equation (11) should be supplemented with initial conditions (in particular, they define constants c_n with $n \geq 1$ in the solution (12)). It is physically reasonable to choose them near the hypersurface of Planckian energy density $V(\phi_{\max}) = M_P^4$ corresponding to the cosmological singularity.³ If the universe started its evolution from the single Hubble patch of the nearly Planckian size (with, say, $H_i^2 \sim \frac{1}{2}M_P^2$), one takes

$$P(\phi, N_i) = \delta\left(\phi - \frac{1}{\sqrt{2}}\phi_{\max}\right), \quad (15)$$

where $N_i = \log a_i$ and a_i is the initial value of the scale factor in that Hubble patch. Observe that one cannot choose initial conditions of the form $P_i = \delta(\phi - \phi_{\max})$ due to breakdown of the quasiclassical approximation used for the derivation of the Langevin equation (9).

From the solution (12) we see that in the deep infrared limit $N = \log a \rightarrow \infty$ or, more precisely, at $\log a \gg E_1^{-1}$ only the term with $E_0 = 0$ survives in the solution (12), i.e., the distribution function $P(\phi, N)$ reaches its time-independent asymptotics

$$P_{LV}(\phi) \sim V(\phi)^{-1} \exp\left(-\frac{3M_P^4}{8V(\phi)}\right) \quad (16)$$

which is called the Linde-Vilenkin wavefunction of the Universe [15, 16]. Therefore, one-point correlation functions of the inflaton field are given in the limit $N \rightarrow \infty$ by the expression

$$\langle \phi^n \rangle_{\text{IR}} = \frac{1}{N} \int_{\phi_{\min}}^{\phi_{\max}} d\phi \phi^n P_{LV}(\phi), \quad (17)$$

where $\phi_{\min}$ is given by the condition $\epsilon(\phi_{\min}) \sim 1$, the normalization is defined as $N = \int_{\phi_{\min}}^{\phi_{\max}} d\phi P(\phi)$, and n is arbitrary (limited only by the condition that the corresponding integrals converge).

Therefore, we have explicitly shown that the naively diverging infrared parts of all one-point correlation functions of the inflaton field are rendered finite once one takes the effects of eternal inflation into account.

One may observe that, although being finite, the correlators $\langle \phi^n \rangle_{\text{IR}}$ still remain very large. As an example, one can take chaotic inflationary model with the potential $V(\phi) = \frac{1}{2}m^2\phi^2$. Using Eqs. (16), (17) and taking into account the the upper bound for the value of the inflaton field is $\phi_{\max} = \sqrt{2}M_P^2/m$, we find that $\langle \phi^2 \rangle_{\text{IR}} \approx 0.96M_P^4/m^2$ and $\langle \phi^4 \rangle_{\text{IR}} \approx 0.15M_P^8/m^4$. The level of non-gaussianity generated by eternal inflation $\langle \phi^4 \rangle_{\text{IR}} - 3\langle \phi^2 \rangle_{\text{IR}}^2 \approx -2.61M_P^8/m^4$ is extremely large. However, this amount of non-gaussianity will never be accessible for an observer within a given Hubble patch since the horizon of this Hubble patch will never cross the hypersurface of eternal inflation [18].

For completeness, let us discuss what happens with the Fokker-Planck probability distribution if eternal inflation comes to an end before the stationary asymptotics (16) is reached.

At $\phi < \phi_{\text{EI}}$ one has to neglect the first, stochastic, term in the right-hand side of the Fokker-Planck equation (11), and the latter acquires the form

$$\frac{\partial P}{\partial N} = \frac{M_P^2}{8\pi} \frac{\partial}{\partial \phi} \left(\frac{1}{V} \frac{\partial V}{\partial \phi} P \right). \quad (18)$$

³ Contrary to the general lore, the regime of eternal inflation does not solve the problem of cosmological singularity, and the latter remains unavoidable even in eternally inflating space-time [17]. By choosing such initial conditions for the Fokker-Planck equation, we neglect Hubble patches where the singularity is achieved but the overall energy density in the patch is dominated by matter fields other than the inflaton field ϕ .

Its solution is given by

$$P(\phi, N) = \frac{V}{V_{,\phi}} f \left(N + \int_{\phi_{\min}}^{\phi} d\phi \frac{8\pi}{M_P^2} V \left(\frac{\partial V}{\partial \phi} \right)^{-1} \right), \quad (19)$$

where the function f should be determined from the evolution of the probability distribution P in the regime of eternal inflation. The physical meaning of the solution (19) is that the probability is transported unchanged along the hypersurface

$$N + \int_{\phi_{\min}}^{\phi} d\phi \frac{8\pi}{M_P^2} V \left(\frac{\partial V}{\partial \phi} \right)^{-1} = N_{\text{total}}. \quad (20)$$

Note that we count the number of e-folds starting from the beginning of inflation instead of from its end, i.e., we define $N = \log(a/a_i)$, where a_i is the initial value of the scale factor.

Finally, we note that the multipoint correlation functions of the form $\langle \phi(x_1)\phi(x_2)\cdots\phi(x_n) \rangle$, where $x_1, x_2, \dots, x_n$ are separated by distances larger than the coarse graining scale (i.e., the horizon scale) depend on the Starobinsky's decoherence parameter δ and cannot be calculated without considering carefully what happens at near-horizon scale.

III. ONE-POINT CORRELATION FUNCTIONS OF THE CURVATURE PERTURBATION

Let us now show that one-point correlation function of the curvature perturbation $\langle \zeta^n \rangle$ also remains finite if one takes properly into account the effects of eternal inflation. Our goal will be to calculate the average duration of the inflationary stage $\langle N_{\text{total}} \rangle$ as well as its higher order correlation functions $\langle N_{\text{total}}^n \rangle$ for arbitrary n . Correlation functions of the curvature perturbation ζ at the end of inflation (i.e., at $N = N_{\text{total}}$) are trivially related to the latter:

$$\langle \zeta^n(N_{\text{total}}) \rangle = \langle (N_{\text{total}} - \langle N_{\text{total}} \rangle)^n \rangle. \quad (21)$$

To calculate these one-point correlation functions, we will use the method developed by Starobinsky in [8] and expand it for the case of chaotic inflationary scenario.

It is basically guaranteed that inflation will come to the end in a given Hubble patch when the inflaton expectation value drops below the boundary ϕ_{EI} defined by the Eq. (7), and the evolution of the inflaton field becomes deterministic. However, there are still stochastic fluctuations of the inflaton field which may lead to a sudden change of ϕ in a given Hubble patch from a value $\phi > \phi_{\text{EI}}$ to the value $\phi_{\min} < \phi \ll \phi_{\text{EI}}$. Thus, to find the total expected number of e-folds for the given Hubble patch one has to use the formalism of stochastic inflation.

The probability distribution for the end of inflation or, in other words, the total number of e-folds N_{total} can be determined from the probability distribution $P(\phi, N)$ from conservation of probability by

$$w(N_{\text{total}}) = P(\phi, N) \left| \left(\frac{\partial \phi}{\partial N_{\text{total}}} \right)_N \right| = \frac{M_P^2}{8\pi} \lim_{\phi \rightarrow \phi_{\min}} \left| \frac{1}{V} \frac{\partial V}{\partial \phi} \right| P(\phi, N_{\text{total}}). \quad (22)$$

Instead of dealing with this distribution function directly, we will calculate the correlation functions

$$Q_n(\phi) = \int_{N_i}^{+\infty} dN N^n P(\phi, N), \quad (23)$$

where $N_i = \log a_i$ and a_i is the value of the scale factor at the beginning of inflation. The moments Q_n can be naturally related to the correlation functions of the total number of e-folds $\langle N_{\text{total}}^n \rangle$ and curvature perturbation correlation functions $\langle \zeta^n \rangle$, as we will see later.

The zero moment $Q_0(\phi)$ can be determined from the equation

$$-\frac{1}{3M_P^2} \frac{\partial^2}{\partial \phi^2} (V Q_0) + \frac{M_P^2}{8\pi} \frac{\partial}{\partial \phi} \left(\frac{1}{V^2} \frac{\partial V}{\partial \phi} (V Q_0) \right) = -P(\phi, N_i). \quad (24)$$

After taking into account that $Q_0(\phi = \phi_{\max}) = 0$, i.e., that the probability flow through the hypersurface of Planckian energy density is absent, one finds

$$V(\phi) Q_0(\phi) = 3M_P^2 e^{-2v(\phi)} \int_{\phi}^{\phi_{\max}} d\phi' e^{2v(\phi')} \int_{\phi_{\min}}^{\phi'} d\phi'' P(\phi, N_i), \quad (25)$$

where $v(\phi)$ is again given by Eq. (13). From this expression one can immediately see that the probability distribution function (22) for the total duration of inflationary stage is properly normalized:

$$\frac{M_P^2}{8\pi} \lim_{\phi \rightarrow \phi_{\min}} \left| \frac{1}{V} \frac{\partial V}{\partial \phi} \right| Q_0 = \int_{N_i}^{\infty} dN w(N) = 1. \quad (26)$$

In turn, the higher moments Q_n are governed by the recursive set of equations

$$-\frac{1}{3M_P^2} \frac{\partial^2}{\partial \phi^2} (V Q_n) + \frac{M_P^2}{8\pi} \frac{\partial}{\partial \phi} \left(\frac{1}{V^2} \frac{\partial V}{\partial \phi} (V Q_n) \right) = -n Q_{n-1}, \quad (27)$$

relating higher moments Q_n with the lower ones Q_{n-1} . The general solution of Eq. (27) can be written in the form similar to (25):

$$V(\phi) Q_n(\phi) = 3M_P^2 e^{-2v(\phi)} \int_{\phi}^{\phi_{\max}} d\phi' e^{2v(\phi')} \int_{\phi_{\min}}^{\phi'} d\phi'' n Q_{n-1}(\phi''). \quad (28)$$

We immediately conclude that all the higher moments $Q_n(\phi)$ are finite, since $Q_0(\phi)$ and all the integrals in this expression are well behaved.

The correlation functions of the total number of e-folds are related with moments $Q_n(\phi)$ according to the prescription

$$\langle N_{\text{total}}^n \rangle = \frac{M_P^2}{8\pi} \lim_{\phi \rightarrow \phi_{\min}} \left| \frac{1}{V} \frac{\partial V}{\partial \phi} \right| Q_n = n \int_{\phi_{\min}}^{\phi_{\max}} d\phi Q_{n-1}(\phi), \quad (29)$$

where we have used Eq. (27).⁴ In particular, the average duration of the inflationary stage is given by

$$\langle N_{\text{total}} \rangle = \int_{\phi_{\min}}^{\phi_{\max}} d\phi Q_0(\phi) = 3M_P^2 \int_{\phi_{\min}}^{\phi_{\max}} d\phi V^{-1}(\phi) e^{-2v(\phi)} \times \quad (30)$$

$$\times \int_{\phi}^{\phi_{\max}} d\phi' e^{2v(\phi')} \int_{\phi_{\min}}^{\phi'} d\phi'' P(\phi, N_i). \quad (31)$$

Finally, the curvature perturbation correlation functions are related to the moments $Q_n(\phi)$ according to the prescription

$$\langle \zeta^n(N_{\text{total}}) \rangle = n \int_{\phi_{\min}}^{\phi_{\max}} d\phi Q_{n-1}(\phi) - \left(\int_{\phi_{\min}}^{\phi_{\max}} d\phi Q_0(\phi) \right)^n. \quad (32)$$

As we see, all one-point correlation functions of N_{total} and ζ (computed at the end of inflationary stage) are finite for all physically interesting chaotic inflationary models. The final answer does depend on the initial conditions for eternal inflation due to the dependence of the zero moment Q_0 on the initial distribution function $P(\phi, N_i)$. For example, one has

$$\langle N_{\text{total}}^n \rangle = C_n \frac{M_P^{2n}}{m^{2n}}, \quad (33)$$

$$\langle \zeta^n(N_{\text{total}}) \rangle = D_n \frac{M_P^{2n}}{m^{2n}} \quad (34)$$

for the $m^2 \phi^2$ chaotic inflation model, where $C_n, D_n = C_n - C_0^n \sim \mathcal{O}(1)$ are numerical constants depending on the initial conditions. By virtue of the definition (21), $D_1 = 0$.

These large non-gaussianities however never become observable for any given observer within a given Hubble patch, and we now turn to a discussion of the observable quantities.

⁴ In the deterministic inflationary regime the first term on the right hand side can be neglected with respect to the second one (see the end of the previous Section).

IV. THE TWO-POINT & THREE-POINT FUNCTIONS OF THE CURVATURE PERTURBATION AT ONE LOOP

In the previous section we demonstrated that the one-point functions of the scalar field and the curvature perturbation are finite when the effects of eternal inflation are taken into account. Thus, there are no real infinities in the theory. Furthermore, the scales alluded to in the previous section are much larger than any currently observable scale. Then the question arises: How do the predictions at scales L translate into predictions at scales $M < L$? We address the issue of relating two different scales in the present section using the prescriptions of the ΔN formalism [2].

According to the ΔN formalism, the perturbative expansion for the correlator of the curvature perturbation to one loop is given by [19]

$$\langle \zeta(\mathbf{x}_1)\zeta(\mathbf{x}_2) \rangle = \left(N_A N^A + N_A N_B^{AB} \langle \phi^2 \rangle_{(L)}^{(l)} \right) G_{12} + \frac{1}{2} N_{AB} N^{AB} G_{12}^2, \quad (35)$$

where $G_{ij} \equiv G(r_{ij}) = G(|\mathbf{x}_i - \mathbf{x}_j|) = \langle \phi(\mathbf{x}_i)\phi(\mathbf{x}_j) \rangle$. The parameters L and l in the symbol $\langle \phi^2 \rangle_{(L)}^{(l)}$ are the infrared and ultraviolet cutoffs, respectively, applied to all quantities in equation (35). The 3-point function to one loop is given by

$$\begin{aligned} \langle \zeta(\mathbf{x}_1)\zeta(\mathbf{x}_2)\zeta(\mathbf{x}_3) \rangle &= \left[N_{AB} N^A N^B + \left(\frac{1}{2} N_{ABC}^A N^B N^C + N_{AB}^A N^{BC} N_C \right) \langle \phi^2 \rangle_{(L)}^{(l)} \right] (G_{12}G_{13} + G_{21}G_{23} + G_{31}G_{32}) \\ &+ \frac{1}{2} N_{ABC} N^{AB} N^C \left((G_{12} + G_{13}) G_{12}G_{13} + (G_{21} + G_{23}) G_{21}G_{23} + (G_{31} + G_{32}) G_{31}G_{32} \right) \\ &+ N_{AB} N^{BC} N_C^A G_{12}G_{13}G_{23}, \end{aligned} \quad (36)$$

where both of the above formulae assume that

$$\langle \phi^A(\mathbf{x}_i)\phi^B(\mathbf{x}_j) \rangle = \delta^{AB} G_{ij}. \quad (37)$$

Thus, $G(r)$ is the correlation function of a generic light field in quasi de-Sitter. The exact scalar field correlator $G(r)$ is supposed to be known up to a superlarge scale L and is presumably well defined – see previous section for its coincident limit. According to the ΔN prescription, all the derivatives of N in (35), (36) are to be evaluated at the value of the background field corresponding to some time after the shortest scale of interest has left the horizon. Note that the expansion (35) requires the existence of a well defined background evolution. Then, the picture of stochastic inflation imposes that the largest field value for which the expansion (35) is applicable is given by the hypersurface of eternal inflation (7) with ϕ_{EI} . For larger field values, $\phi > \phi_{\text{EI}}$, the notion of the background field breaks down and so does the expansion (35). Spacetime above this scale can no longer be described by a global FRW metric and the correlators $G(r)$ are no longer translationally invariant. However, it is important to note that an observer will never interact with the surface of eternal inflation since such an interaction presupposes a universe which has thermalized as only in the thermalized patches will the horizon grow, permitting such measurements.

Consider now an observer located in some random position. Since the state of the field is translationally invariant, the stochastic properties of the field will be described by the correlation function $G(r)$ for any such observer. However, the field fluctuations will not be experimentally accessible beyond some length scale M defining the causal patch of the observer. Fluctuations with wavelengths longer than M should appear as part of the background for this patch. Since these fluctuations only contribute a constant to $G(r)$ for $r < M$, the correlator appropriate for describing measurements within the observer's patch is

$$\tilde{G}(r) \equiv G(r) - \langle \phi^2 \rangle_{(L)}^{(M)}. \quad (38)$$

In other words, an observer limited to measurements in a volume $r < M$ will be able to measure $G(r)$ up to a constant. One has

$$\tilde{G}(r) = \int_{r/L}^{r/M} \frac{dp}{p} \mathcal{P}_\phi(p) \left(\frac{\sin p}{p} - 1 \right) + \int_{1/M}^{1/l} \frac{dk}{k} \mathcal{P}_\phi(k) \frac{\sin kr}{kr} \Rightarrow \tilde{G}(0) = \langle \phi^2 \rangle_{(M)}^{(l)}, \quad (39)$$

where $\mathcal{P}_\phi(k)$ is the power-spectrum obtained from the fourier transform of $G(r)$. The first term in $\tilde{G}$ (39) gives a small finite correction to the correlation function which goes to zero as r becomes much smaller than M . The minimal length scale l corresponds to the shortest cosmological scale of interest; it is known to be at least 60 e-folds smaller than the size of the causal patch M (or larger depending on the temperature at reheating).

Let us now express Eq. (35) in terms of $\tilde{G}(r)$, the correlator appropriate for the patch of the size M . Keeping only one loop terms, we have

$$\langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle = \left(N_A N^A + \frac{1}{2} (N_A N^A)_B^B \langle \phi^2 \rangle_{(L)}^{(M)} + N_A N_B^{AB} \langle \phi^2 \rangle_{(M)}^{(l)} \right) \tilde{G}_{12} + \frac{1}{2} N_{AB} N^{AB} \tilde{G}_{12}^2 + N_A N^A \langle \phi^2 \rangle_{(L)}^{(M)}. \quad (40)$$

The last term is a constant, which cancels out when comparing differences of two correlators. The coefficient of $\tilde{G}(r)$ now includes the term

$$\left\langle N^A N_A \left(\bar{\phi}_L + \phi(\mathbf{x})_{(L)}^{(M)} \right) \right\rangle = N_A N^A \Big|_{\bar{\phi}_L} + \frac{1}{2} (N_A N^A)_B^B \Big|_{\bar{\phi}_L} \langle \phi^2 \rangle_{(L)}^{(M)}, \quad (41)$$

where $\bar{\phi}_L$ denotes the background field appropriate for the scale L and $\phi(\mathbf{x})_{(L)}^{(M)}$ are fluctuations with wavelengths larger than M . The 3-point function gives a similar result. Expressed in terms of $\tilde{G}(r)$ it takes the form

$$\begin{aligned} \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \zeta(\mathbf{x}_3) \rangle &= \left[\left\langle N_{AB} N^A N^B \left(\bar{\phi}_L + \phi(\mathbf{x})_{(L)}^{(M)} \right) \right\rangle + \left(\frac{1}{2} N_{ABC}^A N^B N^C + N_{AB}^A N^{BC} N_C \right) \langle \phi^2 \rangle_{(M)}^{(l)} \right] \left(\tilde{G}_{12} \tilde{G}_{13} + \dots \right) \\ &+ \frac{1}{2} N_{ABC} N^{AB} N^C \left(\left(\tilde{G}_{12} + \tilde{G}_{13} \right) \tilde{G}_{12} \tilde{G}_{13} + \dots \right) + N_{AB} N^{BC} N_C^A \tilde{G}_{12} \tilde{G}_{13} \tilde{G}_{23} \\ &+ 2 N_{AB} N^A N^B \langle \phi^2 \rangle_{(L)}^{(M)} \left(\tilde{G}_{12} + \tilde{G}_{13} + \tilde{G}_{23} \right) + \text{const}, \end{aligned} \quad (42)$$

where we again have retained only terms up to one loop. We thus see that the result of $L \rightarrow M$ is the replacement of the tree level coefficients by their averages,

$$\mathcal{A}_n(M) = \left\langle \mathcal{A}_n \left(\bar{\phi}_L + \phi(\mathbf{x})_{(L)}^{(M)} \right) \right\rangle, \quad (43)$$

where $\mathcal{A}_1 = N^A N_A$, $\mathcal{A}_2 = N_{AB} N^A N^B$ etc, as well as the appearance of a disconnected part which for the two point correlator is simply a constant. Otherwise, the functional form of the expansion as given by the rules of [19] remains the same with all integrals now truncated at the new IR cutoff M . The same should hold for all coefficients of all correlators in the ΔN expansion and therefore we write

$$\langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle_M = \left(\langle N_A N^A \rangle + \langle N_A N_B^{AB} \rangle \langle \phi^2 \rangle_{(M)}^{(l)} \right) \tilde{G}(r) + \frac{1}{2} \langle N_{AB} N^{AB} \rangle \tilde{G}^2(r) + \dots, \quad (44)$$

and

$$\begin{aligned} \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \zeta(\mathbf{x}_3) \rangle_M &= \left[\langle N_{AB} N^A N^B \rangle + \left(\frac{1}{2} \langle N_{ABC}^A N^B N^C \rangle + \langle N_{AB}^A N^{BC} N_C \rangle \right) \langle \phi^2 \rangle_{(M)}^{(l)} \right] \left(\tilde{G}_{12} \tilde{G}_{13} + \dots \right) \\ &+ \frac{1}{2} \langle N_{ABC} N^{AB} N^C \rangle \left(\left(\tilde{G}_{12} + \tilde{G}_{13} \right) \tilde{G}_{12} \tilde{G}_{13} + \dots \right) + \langle N_{AB} N^{BC} N_C^A \rangle \tilde{G}_{12} \tilde{G}_{13} \tilde{G}_{23} \\ &+ 2 \langle N_{AB} N^A N^B \rangle \langle \phi^2 \rangle_{(L)}^{(M)} \left(\tilde{G}_{12} + \tilde{G}_{13} + \tilde{G}_{23} \right) + \dots \end{aligned} \quad (45)$$

Thus, the coefficients appropriate for the cutoff M can be calculated by replacing $\bar{\phi}_L \rightarrow \bar{\phi}_L + \phi(\mathbf{x})_{(L)}^{(M)}$ and averaging over all field fluctuations $\phi(\mathbf{x})_{(L)}^{(M)}$ with wavelengths longer than M . This appears to be the generic form for the ΔN expansion when used in patches smaller than L_{EI} .

Equation (43) could also be interpreted in the following way: for an observer in the patch of the size M making observations at $r < M$, the quantity $\bar{\phi}_L + \phi(\mathbf{x})_{(L)}^{(M)}$ is a slowly varying function, indistinguishable from a background field. Of course, this background differs according to the location of the patch in the larger volume of size M . Thus, the average in equation (43) can be seen as an average over the position of the patch of the size M in the larger volume or, equivalently, an average over all possible “backgrounds” seen by observers residing in M [3, 6]. Of course, this quantity is not (necessarily) related to observable quantities of interest to any particular observer in a specific patch, in agreement with the results of [3, 6].

The above considerations might seem to suggest that the only appropriate infrared cutoff for observers like us is the current horizon $L = 1/H_0$ [3]. Defining the theory at $L = 1/H_0$, seems to avoid the need of using averages in the ΔN expansion. However, framed in this way, the choice of the IR cutoff seems dictated by convenience rather than the theory itself and is hence ad hoc, designed to hide the long wavelength fluctuations. However, fluctuations on scales

larger than our current horizon presumably exist and we “simply” have to wait a few billion years to access them. In the future we will be forced to calculate using the cutoff appropriate for that time and thus obtain a different result for our local curvature perturbation. Therefore, *using a larger box should not introduce any theoretical uncertainty for predictions of the local curvature perturbation field.*

Equation (40) allows us to deal with a change in the IR cutoff. We can define the coefficients in the expansion (35) to be evaluated at a background field appropriate for our current patch and impose $1/H_0$ as an IR cutoff. If now we are granted access to longer wavelengths, say by waiting long enough and repeating our observations, it is expected that we will infer a *different* background evolution since we will now be averaging over a different volume and more long wavelength fluctuations will be visible. Thus, we must accordingly revise the inflationary history that we match to the post-inflationary universe. This will of course lead to a variation in the coefficients of the ΔN expansion for the connected part of the correlators, given by

$$M \frac{d\mathcal{A}_n}{dM} = -\frac{1}{2} (\mathcal{A}_n)_B^B \mathcal{P}_\phi(1/M). \quad (46)$$

This running shows how the connected parts of the correlation functions of the curvature perturbation change as the cutoff scale M is increased. It is akin to the renormalization group flow in quantum field theory, where the coupling constants run with the renormalization scale, and we will accordingly refer to M as the renormalization scale for cosmological perturbations. We note, however, that at least for the 2-point function (40) the renormalization simply amounts to a constant change in the numerical value of the connected part of the correlator (see (35)). The difference of the 2-point functions, which is a measurable quantity, thus remains invariant under the change of the cutoff scale

$$\langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle - \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_3) \rangle = \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle_M - \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_3) \rangle_M. \quad (47)$$

Equation (46) admits the following interpretation, at least in the case of single field inflation. The coefficients $\mathcal{A}_n$ are functions of the derivatives of the number of e-folds $N(\phi)$ w.r.t. ϕ : $\mathcal{A}_n(N', N'', \dots, N^{(n)})$. We assume that from (46) one can derive a consistent set of equations for the running of the derivatives of the form

$$\frac{dN^{(n)}}{d \ln M} = \mathcal{F}(N', N'', \dots, N^{(n)}). \quad (48)$$

We see that as M is changed, the values of the derivatives $N^{(n)}(\phi)$, and thus the function $N(\phi)$ itself, are changed according to (48). Furthermore, equation (48) takes a particularly simple form for inflationary models with a monomial potential $V(\phi) = \kappa \phi^n$. Indeed, for such models

$$N' = -\frac{1}{n} \frac{8\pi}{M_{\text{P}}^2} \phi, \quad N'' = -\frac{1}{n} \frac{8\pi}{M_{\text{P}}^2}, \quad (49)$$

and eqs (48) simply become

$$\frac{dN'}{d \ln M} = -\frac{1}{2} \frac{N''^2}{(N')} \mathcal{P}(1/M). \quad (50)$$

We illustrate this point with a simple example in the next section and refer to future work for further investigation.

V. A CONCRETE EXAMPLE: FREE FIELD

In this section, we consider two point correlators of the curvature perturbation in single field inflation with quadratic potential $V = \frac{1}{2} m^2 \phi^2$, although the following discussion would proceed practically unchanged for any monomial potential. We have

$$N' = -\frac{4\pi}{M_{\text{P}}^2} \bar{\phi}, \quad N'' = -\frac{4\pi}{M_{\text{P}}^2}, \quad (51)$$

and all the higher order derivatives vanish. The two point correlator (35) is thus written to all orders in loops as

$$\langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle = (N')^2 G_{12} + \frac{1}{2} (N'')^2 G_{12}^2. \quad (52)$$

According to (48), the value of N' will change as the cutoff is increased. We parameterize the variation as

$$N' = -\frac{4\pi}{M_P^2} \bar{\phi}_* + \beta(M), \quad \beta(1/H_0) = 0, \quad (53)$$

where we have assumed that the original cutoff has been set at our present horizon $M_0 = 1/H_0$, such that expressions (51) hold for this cutoff, and $\bar{\phi}_*$ denotes the background field appropriate for this cutoff. We thus have

$$\frac{d\beta}{d \ln M} = \frac{1}{2} \left(\frac{4\pi}{M_P^2} \right)^2 \frac{\mathcal{P}(1/M)}{\frac{4\pi}{M_P^2} \bar{\phi}_* + \beta(M)}, \quad (54)$$

from which we obtain

$$\beta(M) = \frac{4\pi}{M_P^2} \bar{\phi}_* \left(1 - \sqrt{1 - \frac{\mathcal{P}}{2\bar{\phi}_*^2} \ln H_0 M} \right) \quad (55)$$

where we have neglected the scale dependence of $\mathcal{P}$. Thus, with the new cutoff M , N' reads

$$N' = -\frac{4\pi}{M_P^2} \bar{\phi}_* \sqrt{1 - \frac{\mathcal{P}}{2\bar{\phi}_*^2} \ln H_0 M} \equiv -\frac{4\pi}{M_P^2} \bar{\phi}_M. \quad (56)$$

In this case, changing M simply amounts to evaluating expressions (51) at a *new* field value, shifted by

$$\bar{\phi} \rightarrow \bar{\phi} \sqrt{1 - \frac{\mathcal{P}}{2\bar{\phi}^2} \ln H_0 M}. \quad (57)$$

Note that the form of the theory remains unchanged, ie still described by a quadratic potential. This is a general feature of all monomial potentials.

So far, we have shown that one is free to choose the IR cutoff for inflationary perturbations, provided appropriate shifts in the parameters of the theory are performed, see (57). It is also interesting to note that for a free field it is simple to construct manifestly infrared finite expressions containing the one-point correlators of the curvature perturbation. The expression for the two point correlator (52) can be inverted to yield

$$G_{12} = \bar{\phi}^2 \left(-1 + \sqrt{1 + (8\pi^2)^{-1} (M_P/\bar{\phi})^4 \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle} \right), \quad (58)$$

where we have neglected the other unphysical root that would yield perturbations of the same order as the background field $G(r) \sim \bar{\phi}^2$. The correlator $G(r)$ in (58) of course depends logarithmically on the IR cutoff but the difference

$$G(r) - G(r') = \int_{1/L}^{1/l} dk \frac{\mathcal{P}_\phi(k)}{k} \left(\frac{\sin(kr)}{kr} - \frac{\sin(kr')}{kr'} \right) = \int_{1/L}^{1/l} dk \frac{\mathcal{P}_\phi(k)}{k} \left(\frac{k}{6} (r'^2 - r^2) + \dots \right) \quad (59)$$

contains only negative powers of L and is thus finite even in the limit $L \rightarrow \infty$. Thus we can construct a manifestly infrared finite expression

$$G_{12} - G_{13} = \bar{\phi}^2 \left(\sqrt{1 + (8\pi^2)^{-1} (M_P/\bar{\phi})^4 \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle} - \sqrt{1 + (8\pi^2)^{-1} (M_P/\bar{\phi})^4 \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_3) \rangle} \right). \quad (60)$$

By approximating the field correlator as

$$G(r) \approx -\mathcal{P}_\phi \ln \left(\frac{r}{L} \right), \quad (61)$$

one further obtains

$$\sqrt{1 + (8\pi^2)^{-1} (M_P/\bar{\phi})^4 \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle} - \sqrt{1 + (8\pi^2)^{-1} (M_P/\bar{\phi})^4 \langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_3) \rangle} = \frac{\mathcal{P}_\phi}{\bar{\phi}^2} \ln \left(\frac{r_{13}}{r_{12}} \right), \quad (62)$$

where the dependence on the cutoff has completely disappeared. This can be seen as a non-perturbative generalization of the first order result (3), which is rederived by expanding (62) to first order in $(M_P/\bar{\phi})^4 \langle \zeta \zeta \rangle$. Since equation (62) does not depend on the cutoff L at all, it remains unchanged in the renormalization prescription described above and is thus renormalization group invariant to the precision of the approximation (61). Given a measurement of the two point correlator of the curvature perturbation $\langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle$ for one separation of points r_{12} , equation (62) can thus immediately be solved to find an unambiguous prediction for the correlator at any other separation r_{13} .

VI. DISCUSSION

The appearance of IR divergences in cosmological perturbation theory has been addressed by many authors with conclusions that do not always coincide [3, 4, 5, 6, 7]. For example, Lyth [3] has suggested that the appropriate IR cutoff for inflationary calculations is a box slightly larger than the observable universe. He argued that the use of a much larger box leads to problems since one would further need to average over the position of our observable patch in this larger box and that this would introduce theoretical uncertainties. However, framed in this way, the choice of the IR cutoff seems dictated by convenience rather than by the theory itself and is hence ad hoc. On the other hand, the authors of [6] have pointed out that IR effects are indeed irrelevant for the observations but conceded that using a box larger than the horizon would require one to average over the position of our patch leading to a prediction for an averaged, and therefore inherently statistical, power spectrum. In contrast, we do not find any such theoretical uncertainty, nor is there any need to average over the possible embeddings of our observable universe into the box defined by the hypersurface of eternal inflation, since the field correlator is manifestly translation invariant. This due to the fact that the box itself is defined by the end of inflation, i.e. each point has experienced the same fixed number of e-folds.

Using the framework of stochastic inflation, we first showed that for slow roll inflation the fluctuations are in fact finite to all scales up to Planckian energy density⁵. However, fluctuations on such scales are unobservable, and only regions which have thermalized are interesting observationally. We then related the predictions for the curvature perturbation on the largest possible thermalized scales to observations performed in much smaller regions by an appropriate redefinition of the coefficients in the ΔN expansion. As we have discussed, this procedure can be carried out to the case of 3-point functions, and we conjecture that this is so for any n-point correlator.

Our findings show that it is indeed permissible to use our horizon as a cutoff but that there is no uncertainty related to using a larger box. In perturbation theory the difference between the two choices is an additive constant plus a redefinition of the background at which the coefficients of the ΔN expansion (35) should be evaluated. In a sense, the size of the box M represents simply the renormalization point, the redefinitions a renormalization prescription. The largest possible box is defined by the requirement that the field has dropped below the eternal inflation threshold (see Eq. (7)) and has therefore thermalized. Since $G(r)$ is translationally invariant, any observer confined in a patch of size M is able to probe it for $r < M$ and obtain the same answer regardless of where the patch is located.

Thus, in general the correlators of the curvature perturbation were seen to depend on the renormalization scale M . Since the (perturbative) renormalization of the two-point correlator amounts to a constant change in the numerical value of the connected part of the correlator, the difference of the two-point functions actually remains invariant under the change of the cutoff scale. As discussed in section V, in the free single field case it is straightforward to find also a non-perturbative generalization of the first order result which is independent of the infrared cutoff L . As is apparent from Eq. (62), given a measurement of the correlation of the curvature perturbation between the points $\mathbf{x}_1$ and $\mathbf{x}_2$, one can find an unambiguous and renormalization point independent prediction for any $\langle \zeta(\mathbf{x}_1)\zeta(\mathbf{x}_3) \rangle$. It would be very interesting to generalize this result to multifield cases as well as to non-trivial potentials.

Acknowledgements

We thank Alexei Starobinsky for illuminating discussions. This work was supported by the EU 6th Framework Marie Curie Research and Training network "UniverseNet" (MRTN-CT-2006-035863) and partly by Academy of Finland grant 114419. S.N. is supported by the GRASPANP Graduate School.

-
- [1] D.N. Spergel, et.al., *Astrophys. J. Suppl.* **170**:377,2007.
 - [2] A. A. Starobinsky *Pis. Zh. Eksp. Teor.* **42**, 124 (1985); A. A. Starobinsky *JETP Lett.* **42**, 124 (1985); M. Sasaki and E. D. Stewart, *Prog. Theor. Phys.* **95**, 71 (1996) [arXiv:astro-ph/9507001]; D. H. Lyth, K. A. Malik and M. Sasaki, *JCAP* **0505**, 004 (2005) [arXiv:astro-ph/0411220]; D. H. Lyth and Y. Rodriguez, *Phys. Rev. Lett.* **95**, 121302 (2005), [arXiv:astro-ph/0504045].
 - [3] D.H. Lyth, *JCAP* **12**, 016 (2007), [arXiv:0707.0361 [astro-ph]].

⁵ While this paper was nearing completion, [7] appeared on the arXiv in which stochastic inflation was used to demonstrate that the one-point function is finite for a special case of $\lambda\phi^4$ theory in pure de-Sitter.

- [4] L. Boubekeur and D. H. Lyth, Phys. Rev. D **73** (2006) 021301 [arXiv:astro-ph/0504046].
- [5] D. H. Lyth, JCAP **0606** (2006) 015 [arXiv:astro-ph/0602285].
- [6] N. Bartolo et. al., arXiv:0711.4263 [astro-ph].
- [7] A. Riotto and M.S. Sloth, arXiv:0801.1845 [hep-ph].
- [8] A.A. Starobinsky, in “*Field Theory, Quantum Gravity and Strings*”, eds. H.J. de Vega and N. Sanchez, Lecture Notes in Physics (Springer-Verlag) 246, 107 (1986).
- [9] Y. Nambu and M. Sasaki, Phys. Lett. B **219**, 240 (1989); A.A. Starobinsky and J. Yokoyama, Phys. Rev. D **50**, 6357 (1994) [arXiv:astro-ph/9407016].
- [10] S. J. Rey, Nucl. Phys. B **284**, 706 (1987); K. i. Nakao, Y. Nambu and M. Sasaki, Prog. Theor. Phys. **80**, 1041 (1988).
- [11] D. S. Salopek and J. R. Bond, Phys. Rev. D **43**, 1005 (1991); S. Mollerach, S. Matarrese, A. Ortolan and F. Lucchin, Phys. Rev. D **44**, 1670 (1991).
- [12] K. Enqvist and M. S. Sloth, Nucl. Phys. B **626**, 395 (2002); D.H. Lyth and D. Wands, Phys. Lett. B **524**, 5 (2002); T. Moroi and T. Takahashi, Phys. Lett. B **522**, 215 (2001) [Erratum, ibid. B **539**, 303 (2002)].
- [13] A. D. Linde, Phys. Lett. B **175**, 395 (1986); A. D. Linde, Mod. Phys. Lett. A **1**, 81 (1986).
- [14] R. P. Woodard, Nucl. Phys. Proc. Suppl. **148**, 108 (2005) [arXiv:astro-ph/0502556].
- [15] A. D. Linde, Sov. Phys. JETP **60**, 211 (1984); A. D. Linde, Lett. Nuovo Cimento **39**, 401 (1984).
- [16] A. Vilenkin, Phys. Rev. D **30**, 509 (1984); A. Vilenkin, Phys. Rev. D **33**, 3560 (1986).
- [17] A. Borde and A. Vilenkin, Phys. Rev. Lett. **72**, 3305 (1994).
- [18] A. D. Linde, “*Particle Physics and Inflationary Cosmology*,” Chur, Switzerland: Harwood (1990) [arXiv:hep-th/0503203].
- [19] C. T. Byrnes, K. Koyama, M. Sasaki and D. Wands, JCAP **0711**, 027 (2007) [arXiv:0705.4096 [hep-th]].